



VERIFICATION OF THE TRANSLATION

I, Won-Jin HONG of 648-23, Yoksam-dong, Kangnam-ku, Seoul, Korea being fully conversant with both the Korean and the English languages, declare and swear that the attached is a true and accurate translation of Korean Patent Application No. 1999-46795 filed on October 27, 1999.

Signed by Won-Jin HONG
Won-Jin HONG

Dated this 7th day of July, 2003

[ABSTRACT]

[Abstract]

The present invention provides sequence listings (SEQ. ID. NO. 1~24) that can be used for the diagnosis and identification of *Mycobacterium tuberculosis* and non-
5 *tubercuolsis Mycobacteria* based on the polymorphism of the genes and provides a method for the diagnosis and identification of the *Mycobacterial* strains. According to the method of the present invention, accurate and prompt diagnosis and identification of *Mycobacterial* strains are accomplished.

[Representing Figure]

Fig. 3

[Key Word]

Mycobacteria tuberculosis, *Mycobacteria*

15

[SPECIFICATION]

[Title of the Invention]

rpoB gene fragment and a method for identifying *Mycobacterium tuberculosis* and
5 *non-tubercuolsis Mycobacterial* strains

[Brief Description of the Drawings]

Fig. 1 is genetic map of *rpoB* gene from *Mycobacterium tuberculosis*.

Fig. 2 is a picture of PCR amplification of *rpoB* gene region (361bp) from the
10 standard stains of *non-tubercuolsis Mycobacteria* (NTM). In this figure, each lane
represents the following strains. M: DNA size marker of 50bp ladder, lane 1: *M.*
gordonae, lane 2: *M. szulgai*, lane 3: *M. kansasii* type I, lane 4: *M. gallinarum*, lane 5: *M.*
avium, lane 6: *M. scrofulaceum*, lane 7: *M. asiaticum*, lane 8: *M. chelonae*, lane 9: *M.*
moriokaese, lane 10: *M. phlei*, lane 11: *M. pulveris*, lane 12: *M. fortuitum* type I, lane 13:
15 *M. austroafricanum*, lane 14: *M. smegmatis*.

Fig. 3 is a picture of electrophoresis of PCR-RFLP (*Msp* I digestion) results of
standard stains of NTM. In this figure, lanes M and 1-14 represent the same strains of Fig.
2, and lane 15 represents *M. marinum*.

Fig. 4 is a picture of electrophoresis of PCR-RFLP (*Hae* III digestion) results of
20 standard stains of NTM. In this figure, lanes M and 1-14 represent the same strains of Fig.
3.

Fig. 5 is a picture of electrophoresis of PCR-RFLP (upper: *Msp* I digestion, lower:
Hae III digestion) result of *M. kansasii* identified by Korean Institute of Tuberculosis. In
this figure lane M represents size marker.

25 Fig. 6 is a picture of electrophoresis of PCR-RFLP (upper: *Msp* I digestion, lower:
Hae III digestion) result of *M. chelonae* identified by Korean Institute of Tuberculosis. In
this figure lane M represents size marker.

Fig. 7 is a picture of electrophoresis of PCR-RFLP (*Msp* I digestion) *M.*
tuberculosis.

30 Fig. 8 is a picture of electrophoresis of PCR-RFLP (*Msp* I digestion) results of *M.*

szulgai, one of the standard stains of NTM.

Fig. 9 shows the comparison of Sequence alignments of the *rpoB* regions from *Mycobacterium tuberculosis* and *non-tubercuolsis Mycobacterial* strains.

5 **[Detailed Description of the Invention]**

[Object of the Invention]

[Technical Field of the Invention and Prior Arts]

The present invention relates to novel genes and to a method for identification and diagnosis of *Mycobacterium tuberculosis* and *non-tubercuolsis Mycobacterial* strains using the genes, more specifically, the present invention relates to *rpoB* gene fragments that have particularities to species of *non-tubercuolsis Mycobacterium* and a method for diagnosing and identifying *Mycobacterium tuberculosis* and *non-tubercuolsis Mycobacterial* strains using the gene fragments.

The genus of Micobacteria comprises *Mycobacterium tuberculosis*, *M. leprae* and other *Micobacteria* species called *non-tuberculosis Mycobacteria* (NTM). Since 1980s, infections of *M. avium* in the AIDS patients have increased throughout the developed countries, and a third of the people in the world are infected and 3 million persons in the world die of them per a year, which cause a great concern to treatment of the tuberculosis.

Timpe and Runyon have classified *Non-tuberculosis Mycobacteria* (NTMs) into 4 classes according to their growth tempo, shape and ability of producing pigment in 1954. That is, they are classified as a photochromogene (I) which grows slowly (generally 5-7 days) and exhibits yellow color when it grows with light; a scotochromogene (II) which exhibits yellow or orange color without light; a nonphotochromogene (III) which exhibits light yellow color or does not exhibit color when it grows with light or without light; and a rapid growers (IV) which grows rapidly compared with other mycobacterium.

However, because various pathologic syndromes regarding mycobacterium are known and new strains are identified and diagnosed, biochemical diagnoses are adopted in addition to the diagnoses using microbiological characteristics. With the microbiological and biochemical diagnoses that are presently adopted, a lot of *Non-tuberculosis Mycobacteria* can be identified and diagnosed with comparably high accuracy. However,

these conventional methods of diagnosis are useful only when the strains form colonies and take long time (8 weeks). And these methods sometimes fail precise identification because when the differences of biochemical test results are trivial, the diagnosis result can be different according to examiners who analyze the results. Therefore, only very skilled expert can obtain reliable diagnosis result, and therefore these methods are not useful for the clinical laboratories.

In addition to the method of identification based on microbiological classification and biochemical test, a method using chromatography to analyze the lipids of cell wall from a mycobacterium or using molecular biological method is used or tried for the identification of the *Non-tuberculosis Mycobacteria* in the advanced countries. The chromatography method is thought to be very efficient because its test particularity is very high; however, because this method needs a step of culturing the strains to obtain samples for the test, the procedures are usually impeded by the slow growth of the strains; and because very expensive equipments are used, its application is restricted to a reference laboratory test only in the advanced countries.

In addition, molecular biological identification method using probes prepared with such genes that show genetic variation only in a same species because maintenance of its sequence listing is high. Because this method uses PCR, this method has such advantages that can amplify target gene with a small number of stains and can shorten the time of identification. A lot of researches using molecular biological method to identify *Mycobacteria* including *Mycobacterium tuberculosis* have been actively studied.

Among these researches, as a method for identifying *Mycobacterium tuberculosis* and *non-tubercuolsis Mycobacterial* strains by using 16S rRNA gene, there is AccuProbe manufactured by Genprobe Company. This method uses rRNA:marked-DNA hybridization and has an advantage that it takes only two days to identify strains; however, because sequence listing maintenance of 16S rRNA gene is very high, sensitivity of identification to the *non-tuberculosis Mycobacteria* is low and respective kits are needed to identify different *Mycobacteria* strains(4-5). As AccuProbe products, kits for five strains of *M. tuberculosis*, *M. avium*, *M. intracellulare*, *M. gordonae*, *M. kansasii* are known, but only one strain is identified in a single experiment using these kits.

So, it is necessary to find a gene fraction having better identification character for *Mycobacterial* species than 16S rRNA and to find a method for identifying multiple strains in a single experiment, and therefore there are a lot of studies to find an improved gene fraction. As a result, *rpoB* gene fraction is thought to be a better gene fraction to identify *Mycobacterial* species than 16S rRNA. A precise identification of strains may be accomplished by amplifying the gene fraction with PCR then analyzing the bases of the PCR products; in addition, it would be possible to identify *Mycobacterium tuberculosis* and *non-tubercuolsis Mycobacterial* strains by using a specific restriction enzyme. Direct sequence analysis of PCR product requires expensive molecular biological equipments and highly skilled technique; therefore this method is difficult to be applied to general clinical laboratories in a general use.

Instead of analyzing the bases of the PCR products, restriction fragment length polymorphism (RFLP) analysis method, which digests the PCR products with restriction enzyme such as *BestEII*, *HaeIII*, *MspI*, *HinfI*, *BsaHI* or the like then compares the size and numbers of the digestion cuts to distinguish generic polymorphisms of the amplified genes so as to finally identify the *Mycobacterial* species, has been invented. This method is simpler than the method of analyzing the bases of PCR products, but has limitations that the genetic polymorphisms should be various enough to distinguish the *Mycobacterial* species in the different species and the genes should be maintained in the same species.

As the gene fractions that satisfy the above condition, *dnaJ* gene, 16S-23S rRNA spacer region 65kDa protein genetic portions are known. The PCR-RFLP method is very simple and more likely to be applied to general clinical laboratories compared with the method of analyzing the sequence listings of PCR products in viewpoint of experiment techniques. However, conventional methods require more than two restriction enzymes to obtain accurate identification; and because the difference of the sizes of gene fractions are very slight, very expensive computers and software in addition to skilled expert to analyze the gene fractions. These difficulties are originated from the facts that the polymorphisms of the target genes used in the PCR-RFLP are not sufficient to distinguish the species.

[Technical Problems that the Present Invention Is To Solve]

A gene fraction, which exists all the *Mycobacterial* species, and whose genetic conservitivity in species is high enough to identify a specific *Mycobacterium* by genetic homogeneity and whose genetic polymorphisms between species are various enough to be applied to molecular biological analysis such as PRA and PCR-DNA to distinguish the
5 *Mycobacterial* species, is required in the identification of *Mycobacterial* species.

The present inventors have searched such gene fraction as above, and as a result, found that gene fractions corresponding to 361bp of rpoB gene are very useful to provide a rapid, simple and accurate method for identifying *non-tubercuolsis Mycobacterial* species; and the present invention accomplished based on the above find.

[Constitution and Function of the Present Invention]

The present invention provides DNA fragments including sequence SEQ. ID. NO. 1 to 4 and 6 to 24, which have genetic conservitivity of existing all *Mycobacterial* species and also have genetic polymorphisms distinguished from other species, so that can be used
15 for identifying *non-tubercuolsis Mycobacterial* species.

In addition, the present invention provides a method for identifying and diagnosing *Mycobacterium tuberculosis* and *non-tubercuolsis Mycobacterial* strains.

The present invention provides a method for identifying *Mycobacterial* strains comprising the step of:

20 (1) digesting a DNA fragment which has one of the sequence Seq. ID. NO 1 to 24 with restriction enzyme to obtain DNA fragment length polymorphism pattern;

(2) isolating DNA fragment from microorganism to identify;

(3) amplifying said DNA fragment;

(4) digesting said amplified DNA fragment with the same restriction enzyme in
25 step (1);

(5) obtaining DNA fragment length polymorphism pattern from DNA fragment in step (4); and

(6) comparing DNA fragment length polymorphism pattern from step (1) with DNA fragment length polymorphism pattern from step (5).

30 Alternatively, the step (1) can be performed at any step before step (6) of

comparing.

Preferably, said restriction enzymes are enzyme HaeIII or MspI. When identification is not proper by using enzyme HaeIII or MspI, enzyme Sau3A1 or BstEII may be used.

Hereinafter, the present invention described in detail together with experiments and the results thereof, but the scope of the present is restricted thereto.

MATERIALS and METHODS

1. *Mycobacterial* species

Mycobacterial species of *Mycobacterium tuberculosis* and *non-tubercuolsis* *Mycobacterium* are shown in Table 1.

Table 1.

Species	ATCC No.
<i>M. abscessus</i>	
<i>M. asiaticum</i>	25276
<i>M. avium</i>	25291
<i>M. aichiense</i>	27280
<i>M. aurum</i>	23366
<i>M. africanum</i>	25420
<i>M. austroafricanum</i>	
<i>M. bovis</i>	19210
<i>M. bovis</i> BCG	
<i>M. chelonae</i>	35749
<i>M. celatum</i>	51131
<i>M. flavescense</i>	
<i>M. fortuitum</i>	6841
<i>M. gallinarum</i>	
<i>M. gilvum</i>	43909
<i>M. gordonae</i>	14470
<i>M. gastri</i>	15754
<i>M. haemophilum</i>	29548
<i>M. intracellulare</i>	13950
<i>M. kansasii</i>	12678
<i>M. malmoense</i>	29571
<i>M. marinum</i>	927
<i>M. moriokaese</i>	
<i>M. peregrinum</i>	14467
<i>M. phlei</i>	11758

<i>M. pulveris</i>	19981 35799 15755
<i>M. scrofulaceum</i>	
<i>M. szulgai</i>	
<i>M. smegmatis</i>	
<i>M. terrae</i>	19423
<i>M. tuberculosis</i>	
<i>M. ulcerans</i>	
<i>M. xenopi</i>	
<i>Nocardia brasiliensis</i>	
<i>N. nova</i>	
<i>Rhodococcus</i>	

2. DNA preparation and PCR amplification

The strains shown in Table 1 were cultured, then a loopful of bacterial colony was taken to be resuspended into 0.4ml of ddH₂O in a 1.5ml tube, and boiled for 10 minutes or sterilized for 15 minutes. Then, the resultants are centrifuged, if necessary.

The reaction composition for PCR contained 50mM of KCl, 10mM of Tris-HCl (pH 8.3), 1.5mM of MgCl₂, 0.001% of gelatin (w/v), each 200μM of dNTPs, 1.25U of Taq, 10pmole of primer. The samples were denatured at a denaturing temperature of 95°C for 5 min for 1 cycle; then amplified performing a cycle that includes denaturation at 94°C for 1 min, primer annealing at 58°C for 1 min and elongation at 72°C for 1 min for 35 times; and at last, the samples were incubated further at the last elongation temperature of 72°C for 10 minutes. The primer sets used in PCR amplification are as follows.

MOTT_{tpo}-5': 5'-TCAAGGAGAAGCGCTACGA-3' (SEQ. ID. NO. 25)

MOTT_{tpo}-3': 5'-GGATGTTGATCAGGGT CTGC-3' (SEQ. ID. NO. 26)

After the PCR, the amplification results were visualized performing 1.5% agarose gel electrophoresis using 5μl of the samples and the sizes of the PCR product (361 bp) were also checked.

3. PCR-RFLP

After confirming successful PCR to obtain desired product size, 17.5μl of PCR products that correspond to approximately 500ng of DNA were digested in a 20μl of reaction volume using 5U (units) of *Msp* I restriction enzyme and 2μl of 10X reaction

buffer solution. After more than 1 hour of incubation at 37°C, 5 µl of 5X DNA loading buffer was added, and then the samples were loaded into a 4% metaphor agarose gel electrophoresis to analyze the PCR product. As a DNA size marker, 100bp ladder (MBI Fermentas, Amherst, NY) was used. In addition, after performing PCR using *M. bovis* DNA as a positive control, then the *Msp* I digestion product of the PCR product was used as inner control size marker.

4. Cloning of PCR product and sequence analysis.

After performing PCR using ATCC standard strains of each NTMs, PCR products were purified by using a GeneClean (BIO101) from agarose gel and cloned into TOPO-TA cloning vector of TOPO-TA cloning kit (In vitrogen). DNA sequences of the cloned PCR products were analyzed.

RESULTS

1. PCR-RFLP analysis of *rpoB* gene using standard strains

Among the NTM standard strains deposited by Korean Institute of Tuberculosis, 33 strains of *Mycobacteria sp.* (from ATCC) and 3 strains belonging to *Nocardia sp.* and *Rhodococcus* were obtained, and genomes were separated from the DNA, then performed PCR on the *rpoB* region using MOTT_{rp} primer set represented by SEQ. ID. NO. 25 and 26. PCR products were visualized with agarose gel electrophoresis to find that 361 bp regions of the *rpoB* were amplified effectively, as expected (Fig. 2). The above-amplified PCR products of each strain were treated with restriction enzymes *Msp* I (Fig. 3) and *Hae* III (Fig. 4) individually, then DNA fragments were analyzed with 4% metaphor agarose gel electrophoresis to find that different size and numbers of DNA fragments were obtained according the characters of the PCR products.

2. PCR-RFLP analysis of *rpoB* gene using clinical isolates of NTM stains isolated by Korean Institute of Tuberculosis

In order to confirm whether the above PCR-RFLP results are obtained only from the standard strains of ATCC or can be obtained from clinical isolates of NTM stains that

are clinically isolated and whether the PCR-RFLP results are species-specific pattern that are conserved in the clinical isolates belonging to the same species, PCR-RFLP analyses were performed on *rpoB* gene of the strains clinically isolated. As the clinical isolates for PCR-RFLP analyses, 109 strains from 10 species, which are thought to be important in the viewpoint of clinical importance and isolation frequency, were selected from the clinical isolate strains that had been identified by strain-identification team of Korean Institute of Tuberculosis based on the microbiological and biochemical examination results. As described above, PCR was performed, PCR products were visualized, and then DNA were digested with *Msp* I and *Hae* III and finally analyzed with 4% metaphor agarose gel electrophoresis (Fig. 5 to Fig. 8).

Fig. 5 shows the result of electrophoresis obtained by PCR amplifying *rpoB* region of *M. kansasii* using DNA of the clinical strains of *M. kansasii* and MOTT*rpo* primer, and digesting the PCR product with *Msp* I (upper) and *Hae* III (lower), and then performing doing electrophoresis. As can be seen in Fig. 5, all the PCR-RFLP results obtained by using *rpoB* regions of every *M. kansasii* strains are the same. Therefore, it is concluded that sequence listings of *rpoB* regions of every *M. kansasii* strains are conserved and it is possible to identify *M. kansasii* by using the sequence listing of *rpoB* region of the *M. kansasii*.

Fig. 6 shows the examination result obtained by using the clinical strains of *M. chelonae*, which are similar with that of *M. kansasii*. That is, all the PCR-RFLP results obtained by using *rpoB* regions of every *M. chelonae* strains are the same; and therefore, it is concluded that sequence listings of *rpoB* regions of every *M. chelonae* strains are well conserved, which suggests that the sequence listing of *rpoB* region of the *M. kansasii* can be applied to identify *M. chelonae*.

On the other hand, because it is difficult to separate 2 subspecies of *M. chelonae*, that is *M. chelonae chelonae* and *M. chelonae absessus*, with conventional biochemical method, they are identified as a *M. chelonae* complex combining the two subspecies together; however with the result of PCR-RFLP method using *rpoB*, PCR-RFLP patterns of *M. chelonae chelonae* in lane 4 and 11 and *M. chelonae absessus* in the other lane have slight difference, and therefore, the sequence listings of *rpoB* genes of the two subspecies

were thought to be different. As a result of analyzing the sequence listings of *rpoB* genes of the two subspecies, it was found that the two subspecies could be distinguished by using restriction enzyme BstEII.

With the above two experiments, it was found that most of the differences between species were detected by the treatment of restriction enzyme *Msp* I, and that the profiles of DNA fragments treated with *Msp* I were similar in the same species, and so only *Msp* I was applied as restriction enzyme in the next experiment using clinical strains.

Fig. 7 and 8 show several examples of PCR-RFLP analysis results of *rpoB* gene regions of clinical strains of *Mycobacteria* sp., and because restriction enzyme profiles of *Msp* I are similar in the strains of *M. tuberculosis* (Fig. 7) and *M. szulgai* (Fig. 8), which can be applied to identify strains. In addition, other experiments were performed with various NTM clinical strains such as *M. gordonae*, *M. avium*, *M. intracellulare*, *M. fortuitum*, *M. malmoense*, *M. scrofulaceum*, *M. terrae*, *M. bovis*, or the like (not described); and as the result, it was found that the PCR-RFLP analysis of *rpoB* gene regions of clinical strains of NTM can be used for identifying strains.

3. Sequence listing analysis of *rpoB* gene fragments of NTM standard strains having clinical importance.

Among the non-tubercuolsis *Mycobacterium* sp. connected with disease, *M. avium-intracellulare* complex, *M. kansasii*, *M. marinum*, *M. fortuitum*, *M. chelonae*, *M. abscessus* or the like are observed frequently, and *M. malmoense*, *M. asiaticum*, *M. xenopi*, *M. simiae*, *M. scrofulaceum*, *M. nonchromogenicum*, *M. smegmatis*, *M. peregrinum*, *M. szulgai*, *M. haemophilum*, *M. ulcerans* or the like are rarely observed; in addition, *M. terrae*, *M. gordonae* or the like are regarded as non-pathogenic non-tubercuolsis *Mycobacterium* sp. but sometimes observed.

Among the above species, *rpoB* gene regions of 20 species (24 strains including *M. tubercuolsi*; 4 strains of *M. gordonae*) were amplified with PCR and their sequence listings were analyzed. The base lengths of *rpoB* genes of the strains analyzed by the present experiment were 361 bp, which is the size of PCR product, and the sequence listings of newly analyzed 235 bp from *rpoB* gene fragment except already known 126 bp on the C-

terminal are described in SEQ. ID. NO. 1 to 24.

4. Algorithm of NTM identification based on the results of PCR-RFLP and sequence listing analysis.

5 Based on the data of PCR-RFLP profile and sequence listing of PCR products of NTM standard strains and clinical strains using restriction enzyme analysis, an algorithm for identifying NTM strains was constructed (Fig. 9).

As seen in Fig. 9, gene regions that having excellent conservative and that having polymorphism are separated well. Therefore, genes from the region having excellent
10 polymorphism can be use as signature gene to identify NTM strains.

[Effect of the Invention]

The gene and PCR-RFLP method of the present invention have a lot of advantages compared with those of conventional methods. In addition to the advantage that it can
15 save time and efforts of the examiner due to its precise and quick identification compared with conventional molecular or biochemical methods, it can identify the strains accurately. Conventionally, it took much time about several months to obtain the results or some strains were not identified with the conventional biochemical method, but these strains can be identified accurately with the present PCR-RFLP method.

REFERENCES

1. The bacillus tuberculosis and the contagiousness of phthisis. *Bost Med Surg J.* 108:235, 1883.

2. Alvarez E., Tavel E.: Recherches sur le bacille de Lustgarten. *Arch. Physiol. Norm. Pathol.*, 3rd series, 6: 303, 1885.
25

3. Timpe A, Runyon EH: The relationship of "atypical" acid-fast bacteria to human disease: A preliminary report. *J Lab Clin Med.* 44: 202, 1954.

4. Kox LFF, van Leeuwen J, Knijper S, Jansen HM and Kolk AHJ: PCR assay based on DNA coding for 16S rRNA for detection and identification of Mycobacteria in
30 clinical samples. *J Clin Microbiol* 33: 3225-3233, 1995.

5. Sanguinetti M, Posteraro B, Ardito F, Zanetti S, Cingolani A, Sechi L, de Luca A, Ortona L and Fadda G: Routine use of PCR-reverse cross-blot hybridization assay for rapid identification of *Mycobacterium* species growing in liquid media. *J Clin Microbiol* 36: 1530-1533, 1998.

5 6. Park, Shim, Cho, Bai, Kim. 1994. Comparison of various primers to detect *Mycobacterium tuberculosis* with PCR. *Journal of Korean Society for Microbiology*, 29(3): 263-268.

7. Bai. 1992. Rapid identification of *Mycobacterium avium* and *Mycobacterium intracellulare* by the amplification of rRNA sequences. 27(5): 443-448.

10 8. The Korean National Tuberculosis Association. 1965, 1970, 1975, 1980, 1985, 1990. Report for the situation of Tuberculosis in Korea.

9. Bai, G. H., Park, K. S., and Kim, S. J.: Clinically isolated mycobacteria other than *Mycobacterium tuberculosis* from 1980 to 1990 in Korea. *J. Kor. Soc. Micro.* 28(1): 1-5, 1993.

15 10. O'brien, R. J., Geiter, L. J., and Snider, D. E.: The epidemiology of nontuberculous mycobacterial diseases in the United States. Results from a national survey. *Am. Rev. Respir. Dis.* 135: 1007-1014, 1987.

11. Contreras, M. A., Cheung, O. T., Sanders, D. E., and Goldstein, R. S.: Pulmonary infection with nontuberculous mycobacteria. *Am. Rev. Respir. Dis.* 137: 149-152, 1988.

20 12. Rogall, T., Wolters, J., Flohr, T. and Botiger, E. C.: Towards a phylogeny and definition of species at the molecular level within the genus *Mycobacterium*. *Int. J. Syst. Bacteriol.* 40(4): 323-330. 1990.

13. Bai, G. H.: Rapid identification of *Mycobacterium avium* and *Mycobacterium intracellulare* by the amplification of rRNA sequences. *J. Kor. Soc. Micro.* 27(5): 443-448, 1992.

14. Woese, C. R. : Bacterial evolution. *Microbiol. Rev.* 51: 221-271, 1987.

15. Gobel, U. B., Geiser, A., and Stanbridge, E. J. : Oligonucleotide probes complementary to variable regions of ribosomal RNA discriminate between *Mycoplasma* species. *J. Gen. Microbiol.* 133: 1969-1974, 1987.

16. Rossau, R., Van Mechale, E., Ley, J. and van Heuverswijn, H.: Specific *Neisseria gonorrhoeae* DNA-probes derived from ribosomal RNA. *J. Gen. Microbiol.* 135: 1735-1745, 1989.

17. Gail, L. W. and Washington II., J. A.: *Mycobacteria* other than *Mycobacterium tuberculosis* : Review of microbiologic and clinical aspects. *Rev. Inf. Dis.* 9(2): 275-294, 1987.

18. Lee, Z. M., Bai, G. H., and Morris, S. L. : The rapid detection of nontuberculous mycobacteria in stool samples using PCR. Presented at 93rd ASM Meeting. U.S.A., 1994.

19. Runyon, E. M. : Anonymous mycobacteria if pulmonary disease. *Med. Clin. North. Am.* 43: 273-290, 1959.

20. Wayne, L. G., and Kubica, G. P.: Family *Mycobacteriaceae* Chester 1897, 63, p.1436-1459. In P.H.A. Sneath, N.S., Mair, M.E. Sharpe, and J.G.Holt(ed.) Bergey's manual of systemic bacteriology, Vol 2. The Williams 2 Wilkins Co., Baltimore, 1986.

21. Jenkins, P. A.: Lipid analysis for the identification of mycobacteria. *An appraisal. Rev. Infect. Dis.* 3: 382-866, 1981.

22. Collins, D. M. G. W. De Lisle : Restriction endonuclease analysis of various strains of *Mycobacterium paratuberculosis* isolated from cattle. *Am. J. Vet.Ras.* 47: 2226-2229, 1986.

23. Rubina, P., J. T. Kuach, and P. Mounts : Isolation and restriction endonuclease analysis of mucobacterial DNA. *J. Gen. Microbiol.* 132: 541-551, 1986.

24. Garcia, M. J. and E. Tabares : Separation of *Mycobacterium gadium* from the rapidly growing mucobacteria on the basis of DNA homology and restriction endonuclease analysis. *J. Gen. Microbiol.* 132: 2265-2269, 1986.

25. Srackebrandt, E. and J. Smida : The phylogeny of the genus *Mycobacterium* as determined by 16S rRNA sequences, and development of DNA-probes. p. 244-250. *Biology og Acinomycetes*. Tokyo, 1998.

26. Stahl, D. A., and J. W. Urbance : The division between fast and slow-growing species corresponds to the natural relationships among the mucobacteria. *J. Bacteriol.* 192: 116-124, 1990.

27. Pitulle, C., Dorsch, W., Kazda, J., Wolters, J., and Stackebrandt, E. :
Phylogeny of rapidly growing members of the genus *Mycobacterium*. *Int. J. Syst.
Bacteriol.* 42: 337-343, 1992.

28. Wallace, R. J., Tschien, J.A. and Stone, M. S.: Spectrum of disease due to
5 rapidly growing mucobacteria. *Rev. Inf. Dis.* 5: 657-679, 1983.

29. Tsukamura, M., H. Shimoide, A. Kuse : Epidemiologic studies of lung disease
due to mucobacteria other than *Mycobacterium tuberculosis* in Japan. *Rev. Inf. Dis.* 3(5):
997-1007, 1981.

30. Jenkins, P. A. : Mycobacteria in the environment. *J. Appl. Bact. Sym. Suppl.*
10 70: 1375-1415, 1991.

31. Woods, G. L. and T. A. Washington II. : Mycobacteria other than
Mycobacterium tuberculosis : Review of microbiologic and clinical aspects. *Rev. Inf. Dis.*
9(2): 275-294, 1987.

[CLAIMS]

[Claim 1]

A DNA fragment which has one of the sequence Seq. ID. NO 1 to 4 or 6 to 24

5

[Claim 2]

A method for identifying *Mycobacterium* strain comprising the step of:

(1) digesting a DNA fragment which has one of the sequence Seq. ID. NO 1 to 24 with restriction enzyme to obtain DNA fragment length polymorphism pattern ;

10

(2) isolating DNA fragment from microorganism to identify;

(3) amplifying said DNA fragment;

(4) digesting said amplified DNA fragment with the same restriction enzyme in step (1) ;

15

(5) obtaining DNA fragment length polymorphism pattern from DNA fragment in step (4) ; and

(6) comparing DNA fragment length polymorphism pattern from step (1) with DNA fragment length polymorphism pattern from step (5).

[Claim 3]

20

A method of claim 2, wherein said DNA fragment length polymorphism from step (1) and step (5) are characterized as obtaining by electrophoresis.

[Claim 4]

25

A method of claim 3, wherein said restriction enzyme is characterized as *HaeIII*, *MspI*, *Sau3A1* or *BstEII*.

[Claim 5]

30

A method of one of the claims 2 to 4, wherein said *Mycobacteria* strain is characterized as *M. tuberculosis*, *M. avium*, *M. abscessus*, *M. flavescent*, *M. africanum*, *M. bovis*, *M. chelonae*, *M. celatum*, *M. fortuitum*, *M. gordonae*, *M. gastri*, *M. haemophilum*,

M. intraeclulare, *M. kansasii*, *M. malmoense*, *M. marinum*, *M. szulgai*, *M. terrae*, *M. scrofulaceum*, *M. ulcerans* or *M. xenopii*.

[FIGURES]

[Fig. 1]

5 [Fig. 2]

[Fig. 3]

10

[Fig. 4]

[Fig. 5]

15

[Fig. 6]

20 [Fig. 7]

[Fig. 8]

[SEQUENCE LISTINGS]

<110> LEE, Hyeyoung

<120> Genes specific for species of Mycobacteria and a method for diagnosis,
identification and characterization of *M. tuberculosis* and
other Mycobacteria

<130> pa99168

<160> 26

<170> KOPATIN 1.5

<210> 1

<211> 208

<212> DNA

<213> *Mycobacterium gordonae* I

<400> 1

tcaaggagaa gcgctacgac ctggcccggg taggccgcta caaggtaac aagaagctcg	60
gcctgcacgt cggcgatccg atcaccagct ccacgtgac cgaggaagac gtcgtcgcca	120
ccatcgagta cctggtccgc ctgcacgagg gccagcacac gatgaccgtc ccgggcggca	180
ccgaggtgcc ggttgagacc gacgacat	208

<210> 2

<211> 208

<212> DNA

<213> *Mycobacterium gordonae* II

<400> 2

tcaaggagaa gcgctacgac ctggcccggg tgggccgcta caaggtaac aagaagctcg	60
gtctgaacgt cggcaagccg atcaccagct cgacgtgac cgaggaagac gtcgtagcca	120
ccatcgagta cctggtgcgg ctgcacgagg gtcagtcggc gatgacgggt cccggcggcg	180
ccgaggtgcc ggttgagacc gacgacat	208

<210> 3

<211> 208

<212> DNA

5 <213> Mycobacterium gordonae III

<400> 3

tcaaggagaa gcgctacgac ctggcccgtg tcggccgcta caaggtcaac aagaagctcg 60
gcctgcacgt cggcgatccg atcaccagct ccacgctgac cgaagaagac gtcgtcgcca 120
10 ccatcgagta cctggtccgt ctgcacgagg gtcagcacac gatgaccgtt ccgggcggca 180
ccgaggttcc ggtggagacc gacgacat 208

<210> 4

<211> 207

15 <212> DNA

<213> Mycobacterium gordonae IV

<400> 4

tcaaggagaa gcgctacgac ctggcccgtg tcggccgcta caaggtcaac aagaagctgg 60
20 gcctgcatgt cggcgatccg atcaccagct cgacgctgac cgaagaggac gtcgtcgcca 120
ccatcgagta cctggtccgc ctccacgagg gtcagcacac gatgacgttc cgggcgggac 180
cgaggttccg gtggagaccg acgacat 207

<210> 5

25 <211> 208

<212> DNA

<213> Mycobacterium tuberculosis

<400> 5

30 tcaaggagaa gcgctacgac ctggcccgcg tcggtcgcta taaggtcaac aagaagctcg 60

ggctgcatgt cggcgagccc atcacgtgt cgacgtgac cgaagaagac gtcgtggcca 120
ccatcgaata tctggccgc ttgcacgagg gtcagaccac gatgaccgtt ccggcgggcg 180
tcgaggtgcc ggtggaaacc gacgacat 208

5 <210> 6
<211> 208
<212> DNA
<213> Mycobacterium terrae

10 <400> 6
tcaaggagaa gcgctacgac ctggcccgcg tcggtcgcta taaggtaac aagaagctcg 60
ggctgcatgt cggcgagccc atcacgtgt cgacgtgac cgaagaagac gtcgtggcca 120
ccatcgaata tctggccgc ttgcacgagg gtcagaccac gatgaccgtt ccggcgggcg 180
tcgaggtgcc ggtggaaacc gacgacat 208

15
<210> 7
<211> 214
<212> DNA
<213> Mycobacterium chelonae

20
<400> 7
tcaaggagaa gcgctacgac ctggcccgcg tgggccggta caaggtaac aagaagctgg 60
gtcttgccgg tgccaaccgc gctctggtga ctgccaccac gctcaccgag gaagacgtcg 120
tcgccaccat cgggtacctg gtgcgcctgc acgagggcca gaccacgatg accgcccccg 180
25 gcggcctcga ggtcccggtc gaggtcgacg acat 214

<210> 8
<211> 208
<212> DNA
30 <213> Mycobacterium kansasii

<400> 8

tcaaggagaa gcgctacgac ctggcccgtg tcggccgata caaggtcaac aagaagctgg 60
gcctgaacac caatcatccg atcaccacga cgacgtgac cgaagaagac gtcgtcgcca 120
5 ccatcgagta tctggtccgc ctgcacgagg gccaggccac gatgaccgtg ccgggcgggg 180
tcgaggtgcc ggtggaaacc gacgacat 208

<210> 9

<211> 223

10 <212> DNA

<213> Mycobacterium scrofulaceum

<400> 9

tcaaggagaa gcgctacgac ctggcccgcg tcggccgcta caaggtcaac aagaagctgg 60
15 gtcgtcacgc cggcgagccg atcacgtcgt ccacgtgac cgaggaagac gtcgtcgca 120
ccatcgaata cctggtccgg ctgcaccacg cccgtacgga tggccagccc gccgtcatga 180
ctgtccccgg cggcacgag gtgccggtgg agaccgacga cat 223

<210> 10

20 <211> 208

<212> DNA

<213> Mycobacterium ulcerans

<400> 10

25 tcaaggagaa gcgctacgac ctggctcgcg tgggtcggta caaggtcaac aagaagctcg 60
gcctgaacgc cgccagccc atcaccagct cgacgtgac cgaggaagac gtcgtcgcca 120
ccatcgaata cctggtccgc ttgcacgagg gccagaccgc gatgaccgtc ccgggcgggtg 180
tcgaggtgcc ggtcgagacc gacgacat 208

30 <210> 11

<211> 208
 <212> DNA
 <213> Mycobacterium marinum

5 <400> 11
 tcaaggagaa gcgctacgac ctggcccggg tgggccggta caaggtaac aagaagctcg 60
 gcctgaacgc cggccagccc atcaccagct cgacgtgac cgaggaagac gtcgtcgcca 120
 ccatcgaata cctgggtccg ttgcacgagg gccagaccgc gatgaccgct ccgggcgggtg 180
 tcgaggtgcc ggtcgagacc gacgacat 208

10
 <210> 12
 <211> 207
 <212> DNA
 <213> Mycobacterium szulgai

15
 <400> 12
 tcaaggagaa gcgctacgac ctggctcgtt cggccgttac aaggtaaca aaaagctcgg 60
 tctgaacgtc ggcgagccga tcaccagttc gacgtgacc gaagaggatg tcgtcgccac 120
 catcgagtac ctggttcggc tgcacgaggg ccagaccacg atgaccgttc ccggcggcac 180
 20 cgaggtgccg gtggagaccg acgacat 207

<210> 13
 <211> 223
 <212> DNA
 25 <213> Mycobacterium gastri

<400> 13
 tcaaggagaa gcgctacgac ctggcccggg tcggccgcta caaggtaac aagaagctgg 60
 gcctgaacac cgatcatccg atcaccacca cgacgtgac cgaagaagac gtcgtcgcca 120
 30 ccatcgagta cctgggtcgc ctgcaccacg cctctcaggg tggccaggcc cccgttatga 180

ctgtccccgg cggggtcgag gtgccggtgg aaaccgacga cat

223

<210> 14

<211> 214

5 <212> DNA

<213> Mycobacterium malmoeense

<400> 14

tcaaggagaa gcgctacgac ctggccaggg ttggccgta caaggtcaac aagaagctcg

60

10 ggctgccggc ggccgagtcg gccgtacccg cctcgaccac gctgaccgaa gcggatgtcg

120

tcgccacat cgagtacctg gtgcgcctgc acgagggcca ggcaacgatg acggttcccg

180

gcggcgtcga ggtgccggtg gagaccgacg acat

214

<210> 15

15 <211> 208

<212> DNA

<213> Mycobacterium avium

<400> 15

20 tcaaggagaa gcgctacgac ctggcccggg tgggcccgtc caaggtcaac aagaagctcg

60

gctgcacgc cggtagccg atcaccagct cgacgtgac cgaggaagac gtcgtcgcca

120

ccatcgagta cctggtgcgc ctgcacgagg gtcagccac gatgaccgtc cccggcggca

180

tcgaggtgcc ggtggagacc gacgacat

208

25

<210> 16

<211> 208

<212> DNA

<213> Mycobacterium bovis

30

<400> 16
tcaaggagaa gcgctacgac ctggcccgcg tcggtcgcta taaggtaac aagaagctcg 60
ggctgcalgt cggcgagccc atcacgtcgt cgacgtgac cgaagaagac gtcgtggcca 120
ccatcgaata tctggtccgc ttgcacgagg gtcagaccac gatgaccgtt ccgggcggcg 180
5 tcgaggtgcc ggtggaaacc gacgacat 208

<210> 17
<211> 208
<212> DNA
10 <213> Mycobacterium celatum

<400> 17
tcaaggagaa gcgctacgac ctgcgcggg tgggccgcta caaggtaac aagaagctcg 60
gcctgaacac cgcgtcccg atcacgacga ccactctgac cgaagaggac gtcgtcgcca 120
15 ccatcgagta cctggtccgc ctgcacgagg gccacaccac gatgaccgtc ccgggcggag 180
tcgaggtgcc ggtggaaacc gacgacat 208

<210> 18
<211> 211
20 <212> DNA
<213> Mycobacterium flavescens

<400> 18
tcaaggagaa gcgctacgac ctggcccgcg tgggtcggta caaggtaac aagaagctgg 60
25 gcatcaccga gaaccggcc gacacgacct cgaccacgt gaccgaagag gacgtcgtcg 120
ccaccatcga gtacctggtg cggctgcatc agggcgacaa gacgatgacc gtcccgggtg 180
gagtcgaggt gcccgctgag gtcgacgaca t 211

<210> 19
30 <211> 208

<212> DNA

<213> Mycobacterium fortuitum

<400> 19

5 tcaaggagaa gcgctacgac ctggcccgcg tgggccgcta caaggtaac aagaagctgg 60
gcctgaacgc cgccagccg atcacgtcgt cgactctgac cgaggaagac gtcgtcgcca 120
ccatcgagta cctggtgcgc ctgcacgagg gccagaccac gatgaccgtc cccggcggcg 180
tcgaggtccc ggtcgaggtg gacgacat 208

10 <210> 20

<211> 205

<212> DNA

<213> Mycobacterium intracellulare

15 <400> 20

tcaaggagaa cgcgtacgac ctggcgcgtg tcggccgcta caaggtaac aagaagctcg 60
gcctgcacgc gggcgagccg atcaccagct cgacgtgac cgaggaagac gtcgtcgcca 120
ccatcgagta cctggtgcgc ctgcacgagg gccagcccac gatgaccgtc cccggcatcg 180
aggcgccggt ggagaccgac gacat 205

20

<210> 21

<211> 214

<212> DNA

25 <213> Mycobacterium abscessus

<400> 21

tcaaggagaa gcgctacgat ctggcccgcg tgggtcggtg caaggtaac aagaagctgg 60
gcctggggcg caccaatccg gtcagggtga ccaccaccac cctcaccgag gaagacgtcg 120
30 tcgccacat cgagtacctg gtgcgcctgc acgagggcca gaccacgatg accgcccccg 180

gcggcgtcga ggtgccggtg gatgtggacg acat

214

<210> 22

5 <211> 208

<212> DNA

<213> Mycobacterium africanum

<400> 22

10 tcaaggagaa gcgctacgac ctggcccgcg tcggtcgcta taaggtaac aagaagctcg 60
ggctcatgt cggcgagccc atcacgtcgt cgacgctgac cgaagaagac gtcgtggcca 120
ccatcgaata tcgtgtccgc ttgcacgagg gtcagaccac gatgatcgtt ccgggcggcg 180
tcgaggtgcc ggtggaaacc gacgacat 208

15

<210> 23

<211> 208

<212> DNA

<213> Mycobacterium haemophilum

20

<400> 23

tcaaggagaa gcgctacgac ctggcccggg ttggtcgta caaggtaac aagaagctcg 60
ggttgacgc cggtagccg atcacgagct cgacgctgac cgaagaggac gtcgtcgcca 120
ccatcgagta cctggtccgg ctgcatgagg gtcagtcgac gatgaccgtt ccaggtggcg 180
25 tcgaggtgcc agtggatact gacgacat 208

<210> 24

<211> 208

<212> DNA

30 <213> Mycobacterium xenopi

<400> 24

tcaaggagaa gcgctacgac ctggcccggg tgggccgcta caaggtcaac aagaaactcg 60
ggctgaacac cgagaatgcg ccaaccacca cgaccctgac cgaagaggac gtcgtcgcca 120
5 ccatcgaata cctgggtgcgc ttgcacgagg ggcacgccac gatgaaggtc cccgggtggcg 180
tcgaggtgcc ggtggagacc gacgacat 208

<210> 25

10 <211> 19

<212> DNA

<213> Artificial Sequence

<220>

15 <223> PCR amplification primer for amplifying the gene of strains of Mycobacterium

<400> 25

tcaaggagaa gcgctacga 19

20 <210> 26

<211> 20

<212> DNA

<213> Artificial Sequence

25 <220>

<223> PCR amplification primer for amplifying the gene of strains of Mycobacterium

<400> 26

ggatgttgat cagggtctgc 20